



**AP<sup>®</sup> CHEMISTRY**  
**2010 SCORING GUIDELINES (Form B)**

**Question 2**  
**(10 points)**



A galvanic cell and the balanced equation for the spontaneous cell reaction are shown above. The two reduction half-reactions for the overall reaction that occurs in the cell are shown in the table below.

Half-Reaction	$E^{\circ}$ (V) at 298 K
$\text{Fe}^{3+}(aq) + e^{-} \rightarrow \text{Fe}^{2+}(aq)$	+0.77
$\text{MnO}_4^{-}(aq) + 8 \text{H}^{+}(aq) + 5 e^{-} \rightarrow \text{Mn}^{2+}(aq) + 4 \text{H}_2\text{O}(l)$	+1.49

(a) On the diagram, clearly label the cathode.

The electrode in the beaker on the right should be labeled.	One point is earned for correct identification of the cathode.
---	--

(b) Calculate the value of the standard potential,  $E^{\circ}$ , for the spontaneous cell reaction.

$E_{\text{cell}} = 1.49 - 0.77 = 0.72 \text{ V}$	One point is earned for the correct numerical answer.
--	---

(c) How many moles of electrons are transferred when 1.0 mol of  $\text{MnO}_4^{-}(aq)$  is consumed in the overall cell reaction?

5.0 moles of electrons are transferred.	One point is earned for the correct numerical answer.
---	---

**AP<sup>®</sup> CHEMISTRY**  
**2010 SCORING GUIDELINES (Form B)**

**Question 2 (continued)**

- (d) Calculate the value of the equilibrium constant,  $K_{eq}$ , for the cell reaction at 25°C. Explain what the magnitude of  $K_{eq}$  tells you about the extent of the reaction.

$\log K_{eq} = \frac{nE}{0.0592} = \frac{5 \times 0.72}{0.0592} = 61$ $K_{eq} = 6.5 \times 10^{60}$ <p>Because the magnitude of <math>K_{eq}</math> is very large, the extent of the cell reaction is also very large and the reaction goes essentially to completion.</p>	<p>One point is earned for the correct substitution.</p> <p>One point is earned for the correct numerical answer.</p> <p>One point is earned for an explanation.</p>
--	--

Three solutions, one containing  $\text{Fe}^{2+}(aq)$ , one containing  $\text{MnO}_4^-(aq)$ , and one containing  $\text{H}^+(aq)$ , are mixed in a beaker and allowed to react. The initial concentrations of the species in the mixture are 0.60 M  $\text{Fe}^{2+}(aq)$ , 0.10 M  $\text{MnO}_4^-(aq)$ , and 1.0 M  $\text{H}^+(aq)$ .

- (e) When the reaction mixture has come to equilibrium, which species has the higher concentration,  $\text{Mn}^{2+}(aq)$  or  $\text{MnO}_4^-(aq)$ ? Explain.

<p><math>[\text{Mn}^{2+}(aq)]</math> will be greater than <math>[\text{MnO}_4^-(aq)]</math> because:</p> <p>(1) as indicated in part (d), the reaction essentially goes to completion, and</p> <p>(2) there is more than sufficient <math>\text{Fe}^{2+}</math> and <math>\text{H}^+</math> to react completely with the <math>\text{MnO}_4^-</math>.</p> <p><math>[\text{MnO}_4^-(aq)]</math> at equilibrium is essentially zero.</p>	<p>One point is earned for the choice of <math>\text{Mn}^{2+}</math> with the explanation including only item (1).</p> <p>One point is earned for including item (2) in the explanation.</p>
--	--

- (f) When the reaction mixture has come to equilibrium, what are the molar concentrations of  $\text{Fe}^{2+}(aq)$  and  $\text{Fe}^{3+}(aq)$ ?

<p>At equilibrium,</p> $[\text{Fe}^{2+}(aq)] = [\text{Fe}^{2+}(aq)]_{\text{initial}} - 5[\text{MnO}_4^-(aq)]_{\text{reacted}}$ $= 0.60 - 5(0.10) = 0.10 \text{ M}$ $[\text{Fe}^{3+}(aq)] = 5 \times [\text{MnO}_4^-(aq)]_{\text{reacted}}$ $= 5(0.10) = 0.50 \text{ M}$	<p>One point is earned for a correct setup (including a correct setup for an equilibrium calculation).</p> <p>One point is earned for correct numerical answers.</p>
---	--